



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Diverging and intergrading forms are abundant in the genus. Noticeably so is it with the arboreal species, *A. canadensis*. Trees occur whose leaves are pubescent throughout the season. Leafy forms occur whose fruit branches are remarkably leafy, the fruit being entirely hidden. Birds make such onslaughts on the ripe fruit that in order to get it in quantity and perfection it is necessary to study it miles away from the inhabited sections, for birds are rare in such localities.

WESTMINSTER, VT.

### SHORTER NOTES

***Ribes chihuahuense*** sp. nov. — Branches smooth, gray. Leaves ovate to suborbicular in outline, the blades 2–2.5 cm. long, dull dark-green above, pale-green beneath, broadly cuneate to subtruncate at the base, 3-lobed, glabrous on both sides, papillose above when young, sparingly ciliate on the margin, the lobes few-toothed, acutish or obtuse, petioles as long as the blades or shorter, pubescent when young; racemes 3–5-flowered, a little longer than the leaves, the axis densely pubescent; flowers sessile or very nearly so, bracts ovate-elliptic, obtuse, ciliate, 5–7 mm. long; hypanthium nearly cylindric, 1 cm. long, sepals oblong, obtuse, 6–7 mm. long; petals ovate-oblong, acute, a little more than half as long as the sepals.

Chihuahua, Mexico, Feb., 1903, *C. A. Purpus*, 1061. Differs from all the United States species by the essentially sessile flowers.

N. L. BRITTON.

FASCIATIONS IN *DROSERA*, *IBERVILLEA*, AND *CECROPIA*. — The fasciated specimen of *Drosera rotundifolia* pictured herewith was found in the propagating houses of the New York Botanical Garden in March, 1907. The flattening affected the stem, resulting in a fasciated rosette, with a growing line 1.4 cm. in length. The literature of teratology seems to contain no instance of fasciation in this genus, while the odd character of the plants makes the appearance of anomalous individuals the object of peculiar interest.

Another fasciation of a curious and rare species is that of a shoot of *Ibervillea Sonorae*. One of the vine-like branches which

spring from the enormous storage tuber measures a little over a centimeter in width. In this case but a single branch is abnormal and the rest of the plant is apparently unaffected.

The fasciation of roots is comparatively infrequent, but Renaudet in a thesis dated Poitiers, 1901, states that it is found oftenest among the aërial roots of tropical species. *Cecropia palmata* produced three fasciated roots in the tropical house of the New York Botanical Garden in 1906. They emerged from the




FIGURE 3. A normal and a fasciated plant of *Drosera rotundifolia*.

trunk somewhat over a foot from the ground, began to flatten close to the main axis, and were finally deeply grooved and bifurcated. The largest measured 12 mm. in width and was three-forked. They resembled closely the drawing of a root of *Pothos aurea* by Udo Dammer in Gardener's Chronicle (26: 724. 1886), although, perhaps owing to the early development of the fasciation, injuries such as he has described in connection with the fasciation of *Pothos* could not be detected.

ALICE ADELAIDE KNOX.